



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## PropiMax<sup>®</sup> EC

### Section 18 Quarantine Exemption

### For Control of Rust (*Phakopsora pachyrhizi*) in Soybean

### For Distribution and Use Only in the State of Michigan

#### ATTENTION

- **This Quarantine Exemption expires November 10, 2007.**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- The product PropiMax<sup>®</sup> EC fungicide, EPA Reg. No. 62719-346, (containing 41.8% propiconazole) manufactured by Dow AgroSciences LLC may be used.
- All applicable conditions and restrictions on the federally registered product label for PropiMax EC as well as the Section 18 use directions must be observed.

#### Active Ingredient:

propiconazole: 1-[[2-(2,4-dichlorophenyl)- 4-propyl-1,3-dioxolan-2-yl]methyl]- 1 <i>H</i> -1,2,4-triazole .....	41.8%
Inert Ingredients .....	58.2%
Total .....	100.0%

Contains petroleum distillates

### Directions for Use

Apply PropiMax EC at the rate of 4 - 8 fl oz per acre (0.113 - 0.225 lb a.i./acre). PropiMax EC may be applied using ground or aerial broadcast equipment or through sprinkler irrigation (chemigation) equipment. Ensure adequate spray volume to achieve good coverage and canopy penetration. For aerial application, apply PropiMax EC in a minimum spray volume of 5 gallons per acre. For best results apply preventatively or as early as possible after an infection has occurred (a delay in fungicide application after an infection period has already occurred may result in yield loss), and make a subsequent application 14 to 21 days later. For maximum residual activity when used as a preventative treatment or for optimum activity on established disease, use the higher rate in the rate range.

**Chemigation:** Apply PropiMax EC in 1/4 to 1/2 inches of sprinkler irrigation. Follow recommendations for application rates, timing and frequency of application in crop-specific directions for use (above).

#### Restrictions:

- A maximum of 3 total applications using approved section 18 products collectively may be made under this soybean rust Section 18 exemption.
- **Do not** make more than 2 applications of propiconazole-containing products to soybeans per season.
- **Do not** apply more than 12 fl oz of PropiMax EC (0.339 lb active ingredient) to soybeans per acre per year.
- **Preharvest Interval:** The final application of PropiMax EC must occur no later than the R5 stage of growth.
- **Re-entry:** A restricted entry interval of 24 hours must be observed.
- **Do not** use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding.
- **Rotational Crop Restrictions:** To avoid possible illegal residues, do not plant any rotational crop intended for food, grazing, or any component of animal feed or bedding within 105 days of application of PropiMax EC to the preceding crop, unless the rotational crop appears on the label for PropiMax EC.

Any adverse effects resulting from the use of PropiMax EC under this emergency exemption must be immediately reported to the Michigan Department of Agriculture.

®Trademark of Dow AgroSciences LLC

R142-057

File symbol: 04-MI-23

EPA-accepted: \_\_/\_\_/\_\_

Replaces R142-023